

Efficacy of Multiple Doses of Antibiotics in Anterior Abdominal Wall Hernia Repair at a Tertiary Rural Hospital

Asadulla Baig¹, Pavan BK², Bhaskaran A^{3*}, Akarsh YG⁴ and Karthik Hareen TVK⁵

¹Assistant Professor, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India

²Assistant Professor, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India

³Professor, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India

⁴Senior Resident, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India

⁵Post-graduate, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India

*Corresponding author: Bhaskaran A, Professor, Department of General Surgery, Sri Devaraj Urs Medical College, Kolar, Karnataka, India, E-mail: dinesh.sathanantham@gmail.com

Received: 15 Mar, 2018 | Accepted: 03 May, 2018 | Published: 09 May, 2018

Citation: Baig A, Pavan BK, Bhaskaran A, Akarsh YG, Karthik Hareen TVK (2018) Efficacy of Multiple Doses of Antibiotics in Anterior Abdominal Wall Hernia Repair at a Tertiary Rural Hospital. J Surg Open Access 4(2): dx.doi.org/10.16966/2470-0991.167

Copyright: © 2018 Baig A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Mesh repair of the anterior abdominal wall hernias is a popular technique and commonly accepted among the majority of surgeons. The technique used most frequently today is the tension free technique. It is uncertain whether extended period of antibiotic prophylaxis is necessary to prevent postoperative wound infection, especially when a foreign body like a polypropylene mesh is used.

Methods: We have studied retrospectively the patients who received surgical treatment in department of Surgery at R L Jalappa Hospital, Tamaka, Kolar for anterior abdominal wall hernia during the period of December 2017- January 2013. Patients were divided into 3 groups based on the doses of antibiotics administered.

Results: In 280 out of 340 cases, a mesh of polypropylene was used. In our sample, we excluded 68 patients due to diseases that made the use of antibiotics necessary. We have studied the frequency of superficial and deep infections in correlation with the use of antibiotics (cephalosporins of third generation).

Conclusion: No difference was observed in the incidence of surgical infection in relation to the duration and the doses of antibiotic cover. The wound infection rate in the current study does not support the use of multiple doses of antibiotics, as this rate does not differ from the rates of infection reported in the literature. Further studies are needed to clarify if antibiotic chemoprophylaxis with one dose or no chemoprophylaxis should be recommended.

Keywords: Hernia; Mesh repair; Wound infection; Chemoprophylaxis

Introduction

The mesh repair of the anterior abdominal wall hernias has been established as the technique of choice among the majority of surgeons. The technique used most frequently today is the free tension technique. This technique allows wound repair, better collagen restoration and prevents recurrence [1]. Furthermore, the use of a polypropylene (PP) mesh has many advantages such as biocompatibility and comfort. This free tension technique is commonly accepted for recurrent, complicated and primary hernias because of the low risk of infection of the introduced foreign body, such as a non-absorbable mesh [2-7]. Surgical wounds may be classified as follows, based on perioperative bacterial contamination: clean, clean contaminated, contaminated and dirty [8-10]. Wound infections are categorized as superficial or deep [8,9,11]. Superficial incisional surgical site infection occurs within 30 days of surgery and involves only the skin and subcutaneous tissue. Deep incisional surgical site infection involves deep soft tissue and appears to be related to the operation. It occurs within 30 days if no implant was left in place and within one year if an implant was left in place [12]. But there are reports of Mesh infection occurring after one year of surgery also.

Hernia mesh repair of the anterior abdominal wall is regarded as a clean surgery and the incidence of postoperative mesh infection is considered to be around 1-2% [1,13,14]. It is uncertain whether antibiotic prophylaxis is necessary to prevent postoperative wound infection, especially when a foreign body like a polypropylene mesh is used. There are no specific guidelines for antibiotic prophylaxis; the surgeon is responsible for determining whether a patient needs antibiotics or not [10,15]. The estimation of the value of the antibiotics seems to be empiric rather than evidence based [4] and that is why their value is a controversial issue. There are antibiotic supportive statements [16-18] that reported

a decrease of the wound infection rate from 9% to 0.7% after antibiotic prophylaxis that are in contrast to other statements in which the antibiotic contribution is underestimated [6].

There is no evidence that the advantages of antibiotic prophylaxis outweigh its disadvantages. The “triple E” [6] summarizes the equivocal nature of antibiotic prophylaxis: ecological impact on the patient’s flora (resistance and mutations of the microorganisms), adverse effects such as anaphylaxis, hypersensitivity, blood dyscrasias and finally, economic impact.

We present our experience with the use of multiple doses of antibiotics in hernia mesh repair.

Purpose of the study

The purpose of this study is to clarify the effectiveness of antibiotic prophylaxis in reducing postoperative wound infection rates in elective open anterior abdominal wall hernia mesh repair.

Materials and Methods

We have studied retrospectively the patients who have undergone surgical treatment in our clinic due to anterior abdomen wall hernia over the past 5 years, from December 2017-January 2013. We have studied the frequency of superficial and deep infections, in conjunction with the use of antibiotics (cephalosporins of third generation or a combination of ampicillin plus sulbactam).

There were 3 groups according to antibiotic prophylaxis duration

Group 1 received antibiotic chemoprophylaxis for 5 days,

Group 2 received antibiotic chemoprophylaxis for 2 days and

Group 3 received 2 doses of antibiotic chemoprophylaxis.

The decisions for allotment of patients were solely based on surgeon’s preference.

In hernia mesh repair of the anterior abdominal wall, we started administering one dose of antibiotic chemoprophylaxis. Comparisons of infection incidence between groups were made using binary logistic regression with group 3 used as a reference category. The confidence interval was 95%, and a difference was considered statistically significant at $p < 0.05$. The statistical analyses were performed using SPSS (Statistical Package for Social Sciences, version 19.0).

The patients were split into 3 groups according to antibiotic prophylaxis duration

Group 1: Of 168 patients who received antibiotic chemoprophylaxis for 5 days, 4 presented with superficial infection and 1 with deep infection.

Group 2: Of 122 patients in who received antibiotic chemoprophylaxis for 2 days, 3 presented with superficial infection and 1 with deep infection.

Group 3: Of 50 patients in who received 2 doses of antibiotic as chemoprophylaxis, 4 presented with superficial infection and 0 with deep infection.

A total of 11 patients presented with superficial infections and 2 patients with deep infections. In both cases, *Staphylococcus aureus* was isolated. The difference in possibility of superficial and deep infection in group 1 compared with group 3 was not statistically significant ($p=0.553$ and 0.995 , respectively) as was the case between groups 2 and 3 ($p=0.887$ and 0.995 , respectively).

Results, Discussion and Conclusions

In our study we did not observe a difference in the infection of the trauma in relation to the duration and the doses of antibiotic cover. Many studies were undertaken to determine the role of antibiotic prophylaxis in mesh hernia repair. An incidence of approximately 10% of the patients submitted to surgical hernia repair presented with infection [4,6,19]. The incidence ranges from 3.3% to 14% (18), and this rate has remained at the same unacceptable level for the past 60 years [14,19-22]. Haley et al. demonstrated a rate of abdominal wall abscess varying from 1.1% to 15.8% [23]. In inguinal herniorrhaphy, surgical site infection is the most frequent complication [24]. It is certain that abdominal wall implant infection increases morbidity. Impaired wound healing and functional loss of the abdominal wall are some of the consequences of implant infection. Also, secondary operations and extended hospital stay are required. A bacterial colonization has been detected in more than 40% of implants [25]. It was demonstrated that bacteria invade the wounds at the time of closure, coming from the body, the air or the surgical instruments [20]. *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Escherichia coli* are usually responsible for mesh infection. Colonies of *Staphylococcus epidermidis* are not vulnerable in prophylactic systemic antibiotics because of a biofilm at the surface of the mesh fibres [25]. Furthermore, the presence of multiple bacterial strains per patient has also been reported [4,16,18,19,26-29]. It has been estimated that 72% of patients are diagnosed during a 4-6 week follow-up period after the surgery, but colonization might occur even after years of implantation and without clinical signs of infection [30]. According to several clinical studies, the occurrence of infection depends on surgical technique and on mesh type. Infection rates, when a polypropylene mesh is used, range from 2-4.2% [31,32]. This infection rate does not differ from the rate observed in the current study. Aufenacker et al. [26] reported a low rate (1.7%) of wound infection after Lichtenstein open mesh inguinal (primary) hernia repair. There was no difference between the antibiotic prophylaxis and the placebo group. Perez et al. [6] also found no difference in the outcome of infection. However, the infection rates are perhaps underestimated due to the unclear definition of infection and the method of follow up. On the other hand, there are several studies that show a significant reduction in infection after antibiotic prophylaxis. Yerdel et al. [18] found that the wound infection rate was 0.7% in the prophylaxis group, and 9% in the placebo group, in addition to the Turkish trial in which the reported infection rates between the group receiving a

single dose of ampicillin plus sulbactam and the placebo group was considerably different. Celdran et al. [7] also reported a reduction in wound infection. However, the sample sizes in those studies were small. Furthermore, a subgroup analysis suggested that in mesh hernia repair, a protective effect could exist and be undetectable because of the small sample size. Furthermore, it has been recently reported that the wound infection rate dropped from 4.2% to 2.3% with the use of prophylaxis [6] and, according to Lazorthes et al. [16], a single-dose of cefandole of 750 mg added to local anesthetic reduced the wound infection rate from 4.5% to 0% compared to anesthesia with no antibiotics [16,18,25]. Troy et al. warranted that a reduction of the growth of bacteria in wounds implanted with mesh was observed after the administration of preoperative single-dose intravenous cefazolin or the topical bacitracin [18]. Gentamycin is reported to be one additional choice for its antimicrobial action. It kills bacteria by diffusing passively across the outer membrane through bacterial pores and, after entering into the cytoplasm, causing the production of a faulty reading of mRNA codons [20]. Gentamycin, in combination with B Lactams [3], results in an antimicrobial synergy and furthermore, at high serum concentrations it can produce auditory and renal damage.

There is research that reported the wound infection rate was at 9% in the placebo group, whereas in the group treated with a mono-dose of ampicillin plus Sulbactam, the infection rate was 0%. Moreover, a single-dose of preoperative intravenous Cefazoline decreases the wound infection rate from 8% to 0%, in comparison to the placebo group [7]. A meta-analysis by Sanabria et al. [33] reported a 50% protective effect of antibiotic prophylaxis on the reduction of wound infection in patients submitted to mesh inguinal hernioplasty. A meta-analysis by Sanchez-Manuel and Seco-Gil [3] for the Cochrane Collaboration reported no statistical difference in SSI rates between antibiotic prophylaxis and no antibiotic prophylaxis groups. The absence of this difference between the teams above was corroborated in the Dutch trial [4,6,18,26].

Moreover, there are studies that examine what antibiotics should be administered and how: intravenously, orally or locally. Terzi et al. [11] reported that a single dose of oral ciprofloxacin is as effective as the administration of intravenous cefazolin in patients undergoing inguinal hernia repair with mesh. Musella et al. [34] reported an infection rate of 0.3% when local antibiotics were used. Several studies reported a remarkable benefit of the intravenous monodose, prophylactic 1.5 gr ampicillin and sulbactam, which resulted in a 3X decrease of the deep incisional surgical site infection and a 10X decrease in overall wound infection. Furthermore, it is certain that proper surgical technique, hemostasis and postoperative surveillance play an important role as far as prevention of wound infection is concerned [8,10]. As for the type of antibiotic, it seems that it is not responsible for the different outcomes of the trials and their use appears to be more important in prosthetic hernia rather than in non-implant repair [6,20]. On the other hand,

in recent reviews Sanchez Manuel et al. [3] concluded that the incidence of superficial wound infection is not altered by the use of a foreign body.

According to recent studies, the drug of choice is ampicillin-clavulanic, which seems to be as effective as cephalosporins. However, this combination has been challenged by a multi-center study [19,23,35], which also supports that there is no difference between oral antibiotic prophylaxis and parenteral drug reception, despite a higher dose of amoxicillin-clavulanic acid being used in the oral treatment group. Furthermore, this combination is twice as expensive as parenteral treatment compared to oral antibiotic prophylaxis.

In summary, the role of antibiotic prophylaxis is still a matter of debate. It is difficult to compare studies due to different antibiotics, surgical techniques, synthetic materials, type of anaesthesia and methods of follow up Sanabria et al. [33] suggest that wound infection rates must be estimated in each hospital to define if antibiotics should be administered in all patients. In case of low rates of wound infection, selective use of antibiotic prophylaxis based on patients' risk factors might be a good option. In addition, cost-effectiveness should be assessed and it should be determined if the benefits of the antibiotic prophylaxis outweigh the drawbacks. Only carefully designed studies are able to answer these questions. Until strong evidence about antibiotics becomes available, surgeons must follow the current guidelines (antibiotics are recommended when there is a high risk of infection or when the occurrence of an infection is associated with severe consequences) [19].

One important step for minimizing infection could be the improvement of the materials [25,36]. A mesh material that uses acrylic acid grafting and gentamicin binding is the antibiotic poly-vinylidene fluoride (PVDF), which was found to have antimicrobial effect with no signs of cell cytotoxicity. A decrease of infiltrating macrophages and apoptotic cells plus physiologic cell proliferation rates is detected by the use of PVDF. It is well understood that the decrease of the infection rate has many benefits in the confinement of the postoperative costs [2] and the complication rate. Regarding the insertion of drains, in general it has been reported that drains acts as a foreign body and may increase the incidence of infection, but the use of drain use in selected patients seems to not increase infection risk [37].

We did not observe a difference in the infection of the surgical site in conjunction with the duration and the doses of antibiotic cover in our 10-year, retrospective study. The wound infection rate in the current study fails to support the use of multiple doses of antibiotics, as the infection rate does not differ from the rates of infection reported in the literature. Today, prosthetic repair of inguinal hernias has low recurrence and infection rates in practice. However, surgical site infection is still a potential complication [38]. Further studies are needed to clarify if antibiotic chemo-prophylaxis with one dose or no chemoprophylaxis should be recommended.

References

1. Engelsman AF, van der Mei HC, Ploeg RJ, Busscher HJ (2007) The phenomenon of infection with abdominal wall reconstruction. *Biomaterials* 28: 2314-2327.
2. Aufenacker TJ, Koelemay MJ, Gouma DJ, Simons MP (2006) Systematic review and meta-analysis of the effectiveness of antibiotic prophylaxis in prevention of wound infection after mesh repair of abdominal wall hernia. *Br J Surg* 93: 5-10.
3. Sanchez-Manuel FJ, Seco-Gil JL (2003) Antibiotic prophylaxis for hernia repair. *Cochrane Database Syst Rev* 2.
4. Tzouvaras G, Delikoukos S, Christodoulides G, Spyridakis M, Mantzos F, et al. (2007) The role of antibiotic prophylaxis in elective tension-free mesh inguinal hernia repair: results of a single-centre prospective randomised trial. *Int J Clin Pract* 61: 236-239.
5. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR (1999) Guideline for Prevention of Surgical Site Infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 27: 97-134.
6. Perez AR, Roxas MF, Hilvano SS (2005) A randomized, double-blind, placebo-controlled trial to determine effectiveness of antibiotic prophylaxis for tension-free mesh herniorrhaphy. *J Am Coll Surg* 200: 393-398.
7. Celdran A, Frieyro O, de la Pinta JC, Souto JL, Esteban J, et al. (2004) The role of antibiotic prophylaxis on wound infection after mesh hernia repair under local anesthesia on an ambulatory basis. *Hernia* 8: 20-22.
8. Biswas S (2005) Elective inguinal hernia repair with mesh: is there a need for antibiotic prophylaxis?--A review. *World J Surg* 29: 830-836.
9. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR (1999) Guideline for prevention of surgical site infection, 1999: Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 20: 279-280.
10. Terzi C (2006) Antimicrobial prophylaxis in clean surgery with special focus on inguinal hernia repair with mesh. *J Hosp Infect* 62: 427-436.
11. Terzi C, Kilic D, Unek T, Hosgorler F, Fuzun M, et al. (2005) Single-dose oral ciprofloxacin compared with single-dose intravenous cefazolin for prophylaxis in inguinal hernia repair: a controlled randomized clinical study. *J Hosp Infect* 60: 340-347.
12. Mann DV, Prout J, Havranek E, Gould S, Darzi A (1998) Late-onset deep prosthetic infection following mesh repair of inguinal hernia. *Am J Surg* 176: 12-14.
13. Deysine M (1998) Pathophysiology, prevention, and management of prosthetic infections in hernia surgery. *Surg Clin North Am* 78: 1105-1115.
14. Taylor SG, O'Dwyer PJ (1999) Chronic groin sepsis following tension-free inguinal hernioplasty. *Br J Surg* 86: 562-565.
15. Knight R, Charbonneau P, Ratzler E, Zeren F, Haun W, et al. (2001) Prophylactic antibiotics are not indicated in clean general surgery cases. *Am J Surg* 182: 682-686.
16. Lazorthes F, Chiotasso P, Massip P, Materre JP, Sarkissian M (1992) Local antibiotic prophylaxis in inguinal hernia repair. *Surg Gynecol Obstet* 175: 569-570.
17. Platt R, Zaleznik DF, Hopkins CC, Dellinger EP, Karchmer AW, et al. (1990) Perioperative antibiotic prophylaxis for herniorrhaphy and breast surgery. *N Engl J Med* 322: 153-160.
18. Yerdel MA, Akin EB, Dolalan S, Turkcapar AG, Pehlivan M, et al. (2001) Effect of single-dose prophylactic ampicillin and sulbactam on wound infection after tension-free inguinal hernia repair with polypropylene mesh: the randomized, double-blind, prospective trial. *Ann Surg* 233: 26-33.
19. Taylor EW, Byrne DJ, Leaper DJ, Karran SJ, Browne MK, et al. (1997) Antibiotic prophylaxis and open groin hernia repair. *World J Surg* 21: 811-814.
20. Deysine M (2006) Infection control in a hernia clinic: 24 year results of aseptic and antiseptic measure implementation in 4,620 "clean cases". *Hernia* 10: 25-29.
21. Medina M, Sillero M, Martinez-Gallego G, Delgado-Rodriguez M (1997) Risk factors of surgical wound infection in patients undergoing herniorrhaphy. *Eur J Surg* 163: 191-198.
22. Santos KR, Bravo Neto GP, Fonseca LS, Gontijo Filho PP (1997) Incidence surveillance of wound infection in hernia surgery during hospitalization and after discharge in a university hospital. *J Hosp Infect* 36: 229-233.
23. Pessaux P, Lermite E, Blezel E, Msika S, Hay JM, et al. (2006) Predictive risk score for infection after inguinal hernia repair. *Am J Surg* 192: 165-171.
24. Bendavid R (1998) Complications of groin hernia surgery. *Surg Clin North Am* 78: 1089-1103.
25. Junge K, Rosch R, Klinge U, Krones C, Klosterhalfen B, et al. (2005) Gentamicin supplementation of polyvinylidene fluoride mesh materials for infection prophylaxis. *Biomaterials* 26: 787-793.
26. Aufenacker TJ, van Geldere D, van Mesdag T, Bossers AN, Dekker B, et al. (2004) The role of antibiotic prophylaxis in prevention of wound infection after Lichtenstein open mesh repair of primary inguinal hernia: a multicenter double-blind randomized controlled trial. *Ann Surg* 240: 955-960.
27. Rios A, Rodriguez JM, Munitiz V, Alcaraz P, Perez Flores D, et al. (2001) Antibiotic prophylaxis in incisional hernia repair using a prosthesis. *Hernia* 5: 148-152.
28. Cruse PJ, Foord R (1980) The epidemiology of wound infection. A 10-year prospective study of 62,939 wounds. *Surg Clin North Am* 60: 27-40.
29. Ehrenkranz NJ (1981) Surgical wound infection occurrence in clean operations; risk stratification for interhospital comparisons. *Am J Med* 70: 909-914.
30. Klosterhalfen B, Klinge U, Hermanns B, Schumpelick V (2000) Pathology of traditional surgical nets for hernia repair after long-term implantation in humans. *Chirurg* 71: 43-51.
31. Lal P, Kajla RK, Chander J, Saha R, Ramteke VK (2003) Randomized controlled study of laparoscopic total extraperitoneal versus open Lichtenstein inguinal hernia repair. *Surg Endosc* 17: 850-856.
32. Neumayer L, Giobbie-Hurder A, Jonasson O, Fitzgibbons R Jr, Dunlop D, et al. (2004) Open mesh versus laparoscopic mesh repair of inguinal hernia. *N Engl J Med* 350: 1819-1827.

33. Sanabria A, Dominguez LC, Valdivieso E, Gomez G (2007) Prophylactic antibiotics for mesh inguinal hernioplasty: a metaanalysis. *Ann Surg* 245: 392-396.
34. Musella M, Guido A, Musella S (2001) Collagen tampons as aminoglycoside carriers to reduce postoperative infection rate in prosthetic repair of groin hernias. *Eur J Surg* 167: 130-132.
35. Norrby SR (1996) Cost-effective prophylaxis of surgical infections. *Pharmacoeconomics* 10: 129-140.
36. Klinge U, Klosterhalfen B, Ottinger AP, Junge K, Schumpelick V (2002) PVDF as a new polymer for the construction of surgical meshes. *Biomaterials* 23: 3487-393.
37. Ergül Z, Akinci M, Yilmaz KB, Sahin A, Seker G, et al. (2011) Why do we use drains in some inguinal hernia repairs? *Chirurgia(Bucur)* 106: 769-774.
38. Genç V, Ensari C, Ergul Z, Kulacoglu H (2010) A very late-onset deep infection after prosthetic inguinal hernia repair. *Chirurgia (Bucur)* 105: 555-557.