

# Laparoscopic Lavage and Drainage in the Management of Complicated Diverticulitis: Review of the Literature

Sainz Hernández Juan Carlos<sup>1\*</sup> and Murillo Zolezzi Adrián<sup>2</sup>

<sup>1</sup>*Surgical Resident, ABC Medical Center, Sur 136 No. 116 Col. Las Américas 01120 Álvaro Obregón, D.F. México*

<sup>2</sup>*Surgeon, ABC Medical Center, Sur 136 No. 116 Col. Las Américas 01120 Álvaro Obregón, D.F. México*

\***Corresponding author:** Juan Carlos Sainz Hernández, Surgical Resident, ABC Medical Center, Sur 136 No. 116 Col. Las Américas 01120 Álvaro Obregón, D.F. México, México, Tel: 52308000, ext. 8661; **E-mail:** [jcsainz@me.com](mailto:jcsainz@me.com)

**Received date:** 09 Sept 2015; **Accepted date:** 12 October 2015; **Published date:** 16 October 2015.

**Citation:** Carlos SHJ, Adrián MZ (2015) Laparoscopic Lavage and Drainage in the Management of Complicated Diverticulitis: Review of the Literature. *J Gastric Disord Ther* 1(2): doi <http://dx.doi.org/10.16966/2381-8689.107>

**Copyright:** © 2015 Carlos SHJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Introduction:** Diverticular disease affects more than 60% of adults aged older than 70. The lifetime prevalence of diverticulitis is 4%-25%, 20% of which will experience severe complications, and 1% will require emergent surgery. We review the most relevant literature regarding the management of complicated acute diverticulitis with laparoscopic lavage and drainage (LLD).

**Discussion:** Hartmann Procedure (HP) is considered the current gold standard for treating complicated diverticulitis; it is associated with a high mortality and morbidity. LLD is reported to improve outcomes. In addition, it may act either as a definitive procedure or as a bridge and allow for a latter elective sigmoidectomy. There is only one preliminary report of a prospective randomized trial regarding this matter (DILALA trial) which fails to demonstrate decreased morbidity and mortality between LLD vs. HP.

**Conclusion:** Longer follow up and results of other trials will be necessary to draw an adequate conclusion.

**Keywords:** Complicated diverticulitis; Laparoscopic lavage; Hartmann's procedure

**Abbreviations:** HP: Hartmann's Procedure; LHP: Laparoscopic Hartmann's Procedure; LLD: Laparoscopic Lavage and Drainage

## Introduction

This paper reviews the current literature on the management of complicated sigmoid diverticulitis: laparoscopic lavage and drainage (LLD) vs. Hartmann's procedure (HP) with the purpose of identifying advantages and disadvantages of each procedure.

## Background

Diverticular disease is common, it affects more than 60% of adults aged older than 70 in the western world [1-3]. Diverticulosis is known as the disease of the industrial revolution, since there is no evidence or pathological reports of this entity prior to the 1900s [4]. In the late 1800s, the process of roller-milling wheat was introduced and it consisted of removing two thirds of the fiber content of wheat. Coincident with this condition, diverticular disease was observed in the first decade of 1900s [4]. It is well known that a low fiber diet contributes to diverticular disease [4-8]. Most patients remain asymptomatic, but acute diverticulitis is a common complication. Diverticulitis is an inflammation of the colon that occurs as a result of the perforation of a diverticulum [5]. The life time prevalence of diverticulitis among patients with diverticulosis has been reported from 4%-25%, some believe this difference may be attributable to changes in diet, rising rates of obesity, and an aging population [9-11].

Among the patients who develop diverticulitis, 15-20% will experience severe complications such a formation of abscess, fistula, obstruction or perforation [12]. Approximately 1% will require emergent surgical intervention [13].

The most commonly used grading system for complicated diverticulitis is the Hinchey classification (Table 1) [9,14]. The management of

diverticulitis depends on the severity and extent of disease. Patients with a localized abscess Hinchey grade I and II are candidates for percutaneous drainage. Most of the patients with generalized peritonitis (Hinchey grade III-IV), have a life-threatening condition which requires emergent surgical management, however the ideal surgical procedure in these situation remains controversial [1,2,10,11,15-18].

## Discussion

In the last 15-20 years the management of perforated diverticulitis has changed tremendously, from Hartmann's procedure described originally for the management of colorectal cancer and later used for complicated diverticulitis [5,18,19], to less aggressive procedures known as "damage control surgery" [8,11,20,21].

Hartmann's procedure has a high rate of procedure-associated mortality and morbidity. In addition, a large number of patients never undergo restitution of intestinal transit (30-84%) [1,2,5,8,11,16-18,21-31]. Percutaneous drainage and LLD make it possible to defer emergent

Hinchey classification	Description
I	Colonic inflammation + Pericolic abscess or phlegmon (confined)
II	Colonic inflammation + Retroperitoneal or pelvic abscess (distant)
III	Colonic inflammation + Purulent peritonitis
IV	Colonic inflammation + Fecal peritonitis

**Table 1:** Hinchey classification

surgical resection in some patients. Both of these procedures allow the peritoneal inflammation to subside and permit a subsequent elective sigmoid resection and primary anastomosis with decreased morbidity and mortality [20]. LLD and drainage has been proposed as an alternative to Hartmann's procedure for Hinchey stage III and IV diverticulitis [10,20,32]. In 1996 O'Sullivan and colleagues (Ireland) [33] described this procedure for acute non-feculent peritonitis. It involves copious washout of the peritoneal cavity and the placement of drains [11,17], with no colonic resection, nor colostomy [1].

Currently, LLD is accepted in the guidelines proposed by the European Association of endoscopic surgeons [29] as well as the Dutch for the management of perforated diverticular disease [11] however it is still not endorsed in the guidelines of the American Society of Colon and Rectal Surgeons (ASCRS).

This procedure, considered a damage control surgery, has shown to have decreased operating time, decreased blood loss and, in some studies, decreased post-operative complications when compared with HP. However, to date there are no prospective randomized controlled trials to provide an adequate level of evidence on which to base a clinical decision. Furthermore, to make the issue more complex, it is unclear what the fate of the patients who undergo LLD is... (Do they require later elective sigmoid resection or observation only?). We will review the most relevant studies addressing this issue. It is important to note the heterogeneity of the studies (including selection criteria) and the bias inherent to the retrospective nature of each of them. Table 2 summarizes the results of the trials reviewed.

Faranda et al. [34] report their experience with LLD for patients with generalized peritonitis that included 16 patients with Hinchey stage III disease and 2 patients with Hinchey stage IV with successful outcomes [35]. The procedure involved spreading biologic fibrin glue directly in the inflamed zone (18 cases), suture repair in 4 cases and omentoplasty in 6 cases. No conversion to laparotomy was needed, there was no mortality and the mean hospital stay was 8 days. Fifteen patients underwent posterior elective laparoscopic sigmoid resection at an interval of 3 to 4 months from the first surgical intervention [31]. This study contrasts sharply with Swank et al. [16] results. They observed 5% mortality and 32% morbidity. Ongoing abdominal sepsis after LLD which required emergent surgery occurred in 13% of the patients. They included only patients with Hinchey stage II and III disease [16].

Liang et al. [17] prospectively compared LLD vs. laparoscopic

Hartmann's procedure (LHP) in 88 consecutive patients; unfortunately they do not provide the selection criteria for either LLD (n=47) or LHP (n=41). They found a statistically significant decrease in operating times (100 vs. 182 min), blood loss (35 vs. 210 ml) and conversion rate (2.1 vs. 14.6%) among the LLD cohort. They report morbidity and mortality together; at 4.3% and 12.5% (only 1 death in the LHP group) it was statistically significant. At long term follow up (time not specified) 44% of the patients in the LLD underwent posterior elective sigmoidectomy, the remaining patients had favorable outcome [17]. In the LHP 72% underwent laparoscopic colostomy closure. The reader of this study must also bear in mind that a LHP in grade III diverticulitis is a complex procedure requiring an experienced laparoscopic surgeon for its completion.

Karoui et al. [24] retrospectively compared LLD vs. laparotomy and primary anastomosis with defunctioning ileostomy (n=35 vs. 24 respectively) in the management of Hinchey III diverticulitis. In the LLD group no conversion to laparotomy was necessary. In the Laparotomy with primary anastomosis no patient required colostomy [23]. No post-operative mortality was reported in either group. When compared to the patients who underwent primary resection and anastomosis with ileostomy post-operative morbidity was higher in the latter group (42%). In this study 26.5% of patients successfully treated by LLD did not undergo further elective sigmoidectomy only one was readmitted 3 weeks after their first event because of recurrent diverticulitis (mean follow up of 21 months). Of the remaining 25 patients all but one underwent elective laparoscopic sigmoid resection [24].

Interestingly Karoui et al. [24] compare the result of those patients who underwent LLD and a later laparoscopic resection (n=25) with those who underwent resection with ileostomy and subsequent ileostomy closure (n=24) and found no mortality in either group. Morbidity was 24% vs. 50% respectively. Although this did not reach statistical significance, it could be attributable to the small sample size. Hospital stay was considerably shorter among the former group (14 vs. 23.5 days in total) [24].

Taylor et al. [18] report a retrospective case series of 14 patients with diverticulitis (2 grade II; 10 grade III; 2 grade IV) initially managed with LLD. 11 patients were discharged without further intervention. 3 remaining patients (2 Hinchey IV; 1 Hinchey III) required reoperation. Eight of these patients underwent further elective sigmoid resection without the need of stoma, 7 of which were performed laparoscopically [18]. Although other authors [34] reported successful outcomes of LLD in feculent peritonitis, Taylor et al. [18] report poor outcomes in these patients.

Author (year of publication)	No. of patients	Hinchey grade LLD II/III/IV	Hinchey grade HP II/III/IV	Morbidity LLD/HP (%)	Mortality LLD/HP (%)	OR time LLD/HP	Blood loss (ml) LLD/HP	Success of LLD (%) (¹)	Hospital stay (days) LLD/HP
Faranda (2000) [34]²	18	0/16/2	NA	16.7/NA	0/NA	NA	NA	NA	8
Taylor (2006) [18]²	14	2/10/2	NA	0/NA	0/NA	NA	NA	11 (79)	6.5
Myers (2008) [37]³	100	25/67/0	0/0/8	11/?	3/?	NA	NA	91	8/18
Swank (2013) [16]²	38	5/29/4	NA	44.7	10.5	68	NA	31 (84)	10
Liang (2011) [17]³,⁴	88	5/36/6	3/31/7	4.3/10.9	0/2.4	<b>99.7/182.9</b>	<b>34.4/210</b>	NA	<b>6.6/16.3</b>
Gentile (2014) [36]²	30	14/2/0	11/3/0	21.4/31.2	7.1/25	<b>75/173</b>	NA	NA	<b>11/19</b>

**Table 2:** Trials summary

¹Success of LLD is defined as control of sepsis, no need for further intervention during same hospital stay.

²Retrospective study

³Prospective study. Only case series, no comparison vs. Hartmann procedure

⁴Laparoscopic Hartmann procedure

⁵NA: Data not available or not recollected

In bold results that are statistically significant

Gentile et al. [36] made a retrospective cohort study that included 30 patients >60 years old with grade II/III diverticulitis. 14 patients underwent LLD and 16 patients open HP. Their analysis favored LLD with improved outcomes in regards to: total operative time, ICU recovery in the early postoperative period, restoration of bowel functions and length of hospital stays (when compared with HP). There was no difference between groups with regards to postoperative morbidity. Although the short-term mortality for LLD vs. HP was 7%, vs. 25%, and mortality at 12 months was 31% vs. 7% respectively, it failed to reach statistical significance. This could be attributable to the small sample size [33]. There was only 62.5% of reversal Hartmann's procedure [36].

Myers et al. [37] prospectively reviewed a case series of 92 patients that underwent LLD for Hinchey II and III diverticulitis. They report morbidity 11% and a mortality of 3%. In this report Myers et al. [37] exclude patients with grade IV diverticulitis, treating them with a HP. Their study mentions the management pathway for their patients, which we agree with, in which grade IV diverticulitis will undergo HP initially, and grade II or III diverticulitis is managed with laparoscopic lavage and drainage with strict in hospital follow up and in whom failure to improve leads to HP.

Rossi et al. [38] retrospectively analyze their data collected from a prospective database in which they included 46 patients who underwent LLD for Hinchey III diverticulitis. In this series 44 of the 46 patients who were chosen to undergo LLD actually underwent LLD (2 were converted). 5 of the 44 patients treated with LLD failed to achieve adequate control of sepsis with the procedure and required further intervention [38].

Cirocchi et al. [39] performed a systematic review in which they conclude that the results from prospective randomized controlled trials are necessary to determine the role of LLD. However, they suggest that LLD can function either as a 1) definitive procedure or 2) as a "bridge" with a later elective sigmoidectomy. The success rate of LLD which they define as patients alive without surgical treatment for recurrent diverticulitis or complications from diverticular disease is 24.3%. They report a 30 day postoperative mortality rate of 4.8% (HP mortality of 19%) [39].

The results of these studies in general favor management of patients with complicated diverticulitis with LLD because of the low morbidity and mortality rates. However, because of their study design the grade of evidence they provide does not support an evidence-based decision. An important bias present in most of the articles regarding this subject is the failure to disclose the selection process by which a patient was taken either to LLD or HP. Moreover, it is not possible converge their results for Meta analysis, as their methodology is far too heterogeneous for comparison.

That said Feingold DL [40] includes in his analysis multiple smaller studies and attempts to make a cumulative analysis of 8 retrospective and 2 prospective case series. This includes 228 patients, 85% were managed with LLD, with a cumulative mortality of 1%. Although long-term follow up is lacking, the author reports a low recurrence rate. In this analysis 24% of the patients had either grade I or II diverticulitis [40].

Various ongoing randomized (Ladies, DILALA, SCANDIV and LapLAND trials) will hopefully shed more light on the issue. Preliminary outcomes of the DILALA trial (comparing LLD with open HP for Hinchey grade III diverticulitis) fail to demonstrate a statistically significant difference in regard to morbidity. Their reoperation rate was 13.2 and 17.1% respectively (P=0.634). They also found no difference in mortality at 30 and 90 days (7.7% vs. 0% (P=0.094), 7.7% vs. 11.4% (P=0.583) respectively). The authors do report a statistically significant decrease in operating time (68 vs. 154 min), and hospital stay (6 vs. 9 days) in the LLD patients. Of the patients treated with LLD non-required reoperation due to ongoing sepsis. The analysis performed is based on a short term follow up, and issues regarding morbidity and mortality of

stoma reversal vs. management of patients treated successfully with LLD are still unaddressed [41].

## Conclusion

Numerous retrospective studies reported regarding LLD and LLD vs. HP seem to be in favor of LLD for the management of grade II/III diverticulitis, with decreased operating time, decreased in hospital stay and, in some reports decreased morbidity. However, because of the limited nature of these trials no valid conclusions can be made. The only prospective randomized trial which to date has been reported fails to show any difference in regards to morbidity and mortality between LLD and HP. Moreover, numerous issues are yet to be resolved: do patients who undergo LLD require elective surgical intervention or follow up? How do the complications of a colostomy and colostomy reversal compare to the long-term results of patients managed with LLD? As more randomized trials become available, we expect to gain further insight as to which surgical strategy offers most benefit.

## References

1. Alamili M, Gögenur I, Rosenberg J (2009) Acute Complicated Diverticulitis Managed by Laparoscopic Lavage. *Dis Colon Rectum* 52: 1345-1349.
2. Tadlock MD, Karamanos E, Skiada D, Inaba K, Talving P, et al. (2013) Emergency surgery for acute diverticulitis: Which operation? A National Surgical Quality Improvement Program study. *J Trauma Acute Care Surg* 74: 1385-1391.
3. Ho VP, Nash GM, Milsom JW, Lee SW (2015) Identification of diverticulitis patients at high risk for recurrence and poor outcomes. *J Trauma Acute Care Surg* 78: 112-119.
4. Painter NS, Burkitt DP (1975) Diverticular disease of the colon, a 20th century problem. *Clin Gastroenterol* 4: 3-21.
5. Moore FA, Catena F, Moore EE, Leppaniemi A, Peitzmann AB (2013) Position paper: management of perforated sigmoid diverticulitis. *World J Emerg Surg* 8: 55.
6. Painter NS (1982) Diverticular disease of the colon. The first of the western diseases shown to be due to a deficiency of dietary fibre. *S Afr Med J* 61: 1016-1020.
7. Unlu C, Daniels L, Vrouenraets BC, Boermeester MA (2012) A systematic review of high-fibre dietary therapy in diverticular disease. *Int J Colorectal Dis* 27: 419-427.
8. Tan KK, Liu JZ, Shen SF, Sim R (2011) Emergency surgery in colonic diverticulitis in an Asian population. *Int J Colorectal Dis* 26: 1045-1050.
9. Bridoux V, Antor M, Schwarz L, Cahais J, Khalil H, et al (2014) Elective operation after acute complicated diverticulitis: Is it still mandatory? *World J Gastroenterol* 20: 8166-8172.
10. Li D, Baxter NN, McLeod RS, Moineddin R, Wilton AS, et al. (2014) Evolving Practice Patterns in the Management of Acute Colonic Diverticulitis: A Population-Based Analysis. *Dis Colon Rectum* 57: 1397-1405.
11. Collins D, Winter DC (2014) Laparoscopy in diverticular disease: Controversies. *Best Pract Res Clin Gastroenterol* 28: 175-182.
12. Parks TG (1969) Natural history of diverticular disease of the colon. A review of 521 cases. *Br Med J* 12: 85-109.
13. Cirocchi R, Trastulli S, Desiderio J, Listorti C, Boselli C, et al. (2013) Treatment of Hinchey stage III-IV diverticulitis: a systematic review and meta-analysis. *Int J Colorectal Dis* 28: 447-457.
14. Wieghard N, Geltzeiler CB, Tsikitis VL (2015) Trends in the surgical management of diverticulitis. *Ann Gastroenterol* 28: 25-30.
15. Letarte F, Hallet J, Drolet S, Charles Grégoire R, Bouchard A, et al. (2013) Laparoscopic Emergency Surgery for Diverticular Disease

- That Failed Medical Treatment: A Valuable Option? Results of a Retrospective Comparative Cohort Study. *Dis Colon Rectum* 56: 1395-1402.
16. Swank HA, Mulder IM, Hoofwijk AG, Nienhuijs SW, Lange JF, et al. (2013) Early experience with laparoscopic lavage for perforated diverticulitis. *Br J Surg* 100: 704-710.
  17. Liang S, Russek K, Franklin ME Jr (2012) Damage control strategy for the management of perforated diverticulitis with generalized peritonitis: laparoscopic lavage and drainage vs. laparoscopic Hartmann's procedure. *Surg Endosc* 26: 2835-2842.
  18. Taylor CJ, Layani L, Ghusn MA, White SI (2006) Perforated diverticulitis managed by laparoscopic lavage. *ANZ J Surg* 76: 962-965.
  19. Boyden AM (1950) The surgical treatment of diverticulitis of the colon. *Ann Surg* 132: 94-109.
  20. Psarras K, Symeonidis NG, Pavlidis ET, Micha A, Baltatzis ME, et al. (2011) Current management of diverticular disease complications. *Tech Coloproctol* 15: S9-S12.
  21. Regenbogen SE, Hardiman KM, Hendren S, Morris AM (2014) Surgery for diverticulitis in the 21st century: a systematic review. *JAMA Surg* 149: 292-302.
  22. Constantinides VA, Heriot A, Remzi F, Darzi A, Senapati A, et al. (2007) Operative Strategies for Diverticular Peritonitis, A Decision Analysis Between Primary Resection and Anastomosis Versus Hartmann's Procedures. *Ann Surg* 245: 94-103.
  23. Afshar S, Kurer MA (2012) Laparoscopic peritoneal lavage for perforated sigmoid diverticulitis. *Colorectal Dis* 14: 135-142.
  24. Karoui M, Champault A, Pautrat K, Valleur P, Cherqui D, et al. (2009) Laparoscopic peritoneal lavage or primary anastomosis with defunctioning stoma for Hinchey 3 complicated diverticulitis: results of a comparative study. *Dis Colon Rectum* 52: 609-615.
  25. Turley RS, Barbas AS, Lidsky ME, Mantyh CR, Migaly J, et al. (2013) Laparoscopic versus open Hartmann procedure for the emergency treatment of diverticulitis: a propensity-matched analysis. *Dis Colon Rectum* 56: 72-82.
  26. Sallinen VJ, Mentula PJ, Leppäniemi AK (2014) Nonoperative Management of Perforated Diverticulitis With Extraluminal Air Is Safe and Effective in Selected Patients. *Dis Colon Rectum* 57: 875-881.
  27. Thornell A, Angenete E, Gonzales E, Heath J, Jess P, et al. (2011) Treatment of acute diverticulitis laparoscopic lavage vs. resection (DILALA): study protocol for a randomised controlled trial. *Trials* 12: 186.
  28. Aydin HN, Tekkis PP, Remzi FH, Constantinides V, Fazio VW (2006) Evaluation of the risk of a non restorative resection for the treatment of diverticular disease: the Cleveland Clinic diverticular disease propensity score. *Dis Colon Rectum* 49: 629-639.
  29. Salem L, Flum DR (2004) Primary anastomosis or Hartmann's procedure for patients with diverticular peritonitis? A systematic review. *Dis Colon Rectum* 47: 1953-1964.
  30. Constantinides VA, Tekkis PP, Athanasiou T, Aziz O, Purkayastha S, et al. (2006) Primary Resection With Anastomosis vs. Hartmann's Procedure in Nonelective Surgery for Acute Colonic Diverticulitis: A Systematic Review. *Dis Colon Rectum* 49: 966-981.
  31. Sauerland S, Agresta F, Bergamaschi R, Borzellino G, Budzynski A, et al. (2006) Laparoscopy for abdominal emergencies: evidence-based guidelines of the European Association for Endoscopic Surgery. *Surg Endosc* 20: 14-29.
  32. Hupfeld L, Burcharth J, Pommergaard H-C, Rosenberg J (2014) The Best Choice of Treatment for Acute Colonic Diverticulitis with Purulent Peritonitis Is Uncertain. *BioMed Research International* 2014: 380607.
  33. O'sullivan GC, Murphy D, O'Brien MG, Ireland A (1996) Laparoscopic management of generalized peritonitis due to perforated colonic diverticula. *Am J Surg* 171: 432-434.
  34. Faranda C, Barrat C, Catheline JM, Champault GG (2000) Two-stage Laparoscopic Management of Generalized Peritonitis Due to Perforated Sigmoid Diverticula: Eighteen Cases. *Surg Laparosc Endosc Percutan Tech* 10: 135-138.
  35. Andersen JC, Bundgaard L, Elbrond H, Laurberg S, Walker LR, et al. (2012) Danish national guidelines for treatment of diverticular disease. *Dan Med J* 59: C4453.
  36. Gentile V, Ferrarese A, Marola S, Surace A, Borello A et al. (2014) Perioperative and postoperative outcomes of perforated diverticulitis Hinchey II and III: Open Hartmann's procedure vs. laparoscopic lavage and drainage in the elderly. *Int J Surg* 12: S86-S89.
  37. Myers E, Hurley M, O'Sullivan GC, Kavanagh D, Wilson I, et al. (2008) Laparoscopic peritoneal lavage for generalized peritonitis due to perforated diverticulitis. *Br J Surg* 95: 97-101.
  38. Rossi GL, Mentz R, Bertone S, Ojea Quintana G, Bilbao S, et al. (2014) Laparoscopic peritoneal lavage for Hinchey III diverticulitis: is it as effective as it is applicable? *Dis Colon Rectum* 57: 1384-1390.
  39. Cirocchi R, Trastulli S, Vettoretto N, Milani D, Cavaliere D, et al. (2015) Laparoscopic peritoneal lavage: a definitive treatment for diverticular peritonitis or a "bridge" to elective laparoscopic sigmoidectomy?: a systematic review. *Medicine (Baltimore)* 94: e334.
  40. Feingold DL (2011) Laparoscopic lavage for hinchey grade III sigmoid diverticulitis. *Semin Colon Rectal Surg* 22: 173-179.
  41. Angenete E, Thornell A, Burcharth J, Pommergaard HC, Skullman S, et al. (2014) Laparoscopic Lavage Is Feasible and Safe for the Treatment of Perforated Diverticulitis With Purulent Peritonitis: The First Results From the Randomized Controlled Trial DILALA. *Ann Surg*.