

Fontan Operation Experience and Results of our Cardiovascular Surgery Center

Farid Gojaye^{1,*}, Ruya Cholak², and Emin Tireli³

¹Department of Cardiovascular Surgery, Bağcılar Training and Research Hospital; Istanbul, Turkey

²Department of Pediatrics, Altınbaş University Medical Park, Bahçelievler Hospital, Istanbul, Turkey

³Department of Cardiovascular Surgery, Altınbaş University Medical Park, Bahçelievler Hospital, Istanbul, Turkey

*Corresponding author: Farid Gojaye, Department of Cardiovascular Surgery, Bağcılar Training and Research Hospital; Istanbul, Turkey; E-mail: farid.gojaye@yahoo.com

Received: 09 Sep, 2022 | Accepted: 27 Oct, 2022 | Published: 02 Nov, 2022

Citation: Gojaye F, Cholak R, Tireli E (2022) Fontan Operation Experience and Results of our Cardiovascular Surgery Center. J Surg Open Access 8(3): dx.doi.org/10.16966/2470-0991.267

Copyright: © 2022 Gojaye F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: In this study, we present our experiences and results of the first Fontan surgeries we performed in our center in accordance with the surgical criteria we believe in.

Materials and methods: This is an observational study involving 19 patients. We investigated the data and survival rate of 19 patients who underwent Fontan surgery one by one. We wanted to know how the surgical method we used and the age factor affected our results. Our main hypothesis was to keep the cardiopulmonary bypass time very short and not to prefer the total circulatory arrest method.

Results: Patients were not homogeneous in terms of demographic characteristics. Patients differed in terms of pre- and postoperative cardiac diagnosis and complications. We lost 2 of 19 patients and the overall mortality rate was 10.5%. We used Extracardiac Membrane Oxygenation (ECMO) support in 3 patients. We applied Implantable Cardioverter Defibrillator (ICD) implantation to the patient. The discharge due to effusion due to chylothorax was prolonged in 4 patients. The postoperative NYHA capacity of the patients was evaluated as 1 or 2.

Discussion: We recommend that this operation should not be performed at a very young age, if the oxygen saturation value of heart patients who are candidates for Fontan is not very critical, if the ventricular ejection fraction capacity is not low, if the physical exertion capacity and examination findings are within normal limits.

Conclusion: The Fontan procedure is still the best long-term palliative operation for all single ventricle cases. We believe that it would be beneficial to postpone this surgery to the advanced age group as much as possible in order not to increase the mortality rate.

Keywords: Fontan procedure; Congenital heart disease; Univentricular heart

Introduction

When performing the surgical treatment of univentricular heart defects, there is a need to create a well-functional single ventricle that will provide balanced circulation to both the lungs and the body. The last step in the surgical correction of univentricular morphological heart defects is the Fontan procedure [1]. The main purpose of this surgery is to direct the circulation of the entire venous system directly to the pulmonary system without contacting the heart and to reduce the excess heart load. Old modification of classical Fontan surgery first performed in 1971 [2]. Until then, many patients born with a single ventricle would usually die within the first year of life. The aim of this multi-stage therapy is to reduce the volume load of the single ventricle by separating the pulmonary circulation from the systemic circulation, and to keep arterial blood saturation values close to normal by balancing the ratio (Qp/Qs) between pulmonary and systemic flow [3]. Thus, we can improve the long-term life span and quality. According to the latest published data, the 30-year survival expected to be approximately 89% (25) [4].

Materials

We performed a total of 279 congenital heart surgeries between 2018-2020. Of these, 19(6.81%) patients reached the Fontan procedure stage and we performed Fontan surgery on these patients. This was an observational study. We investigated the data and survival rate of 19 patients who underwent Fontan surgery one by one. We wanted to know how the surgical method we used and the age factor affected our results. Our main hypothesis was to keep the cardiopulmonary bypass time very short and not to prefer the total circulatory arrest method. Eleven of the patients were male and eight were female. According to the age limitation, 4 people were under 4 years old and 15 people were between 4-8 years old. The patient distribution according to primary cardiac anomalies was 1 Ebstein anomaly, 7 Tricuspid Atresia (TA), 2 tricuspid atresia+Pulmonary Stenosis (PS), 5 Hypoplastic Left Heart Syndrome (HLHS), 2 double-inlet left ventricle, 2 transposition of the great arteries (TGA)+PS. Additional pathologies were Atrial Septal Defect (ASD), Ventricular Septal Defect (VSD). The classification of the patients according to their primary diagnosis is shown in table

Table 1: Distribution of patients included in the study according to their primary diagnosis.

Total Patients	TA	TA+PS	DILV	TGA+PS	HLHS	Ebstein Anomaly
19	7	2	2	2	5	1

*TA: Tricuspid Atresia

*PS: Pulmonary Stenosis

*BAT: Great Artery Transposition

*HLHS: Hypoplastic Left Heart Syndrome

*DILV: Double-Inlet Left Ventricle

1. Left pulmonary artery of a patient with TGA+PS was critically hypoplastic. We completed the Fontan circulation in this patient with the existing unilateral right pulmonary artery (Figure 1). Blalock-Taussig shunt had been applied to three patients previously to improve pulmonary blood circulation. In a study on the complaints of edema, low saturation and bruising in a TA+PS patient 3 months after Glenn shunt surgery, stenosis in the anastomosis line was found. There upon, we applied balloon to the stenotic area and stenting to the right ventricular outflow tract to improve antegrade pulmonary flow (Figure 2). Upon stabilization of the patient's condition, we underwent an elective Fontan operation. The application complaints of all cases consisted of findings such as bruising, shortness of breath, tachypnea, fatigue, decreased effort capacity and palpitation growth and development retardation, and heart failure.

Methods

Antegrade aortic cannulation was performed with a median sternotomy under general anesthesia. Selective cannulation of superior and inferior vena cava was performed in all patients. Mild hypothermia 28-32°C was achieved under Continuous Cardiopulmonary Bypass (CPB). During the cooling, we completed the adherent tissue dissections. Particularly, regions with continuity of pulmonary arteries located behind the ascending aorta were liberated. We used only one dose of hot blood cardioplegia (30 ml/kg) through the aortic antegrade route during the entire operation. We took care to keep the cross-clamp time short (13.2 ± 8.7 min). We did not prefer the total circulatory arrest method in any patient. All cases had previously undergone a bidirectional Glenn shunt operation. We did not use a patch for tricuspid valve closure between the right atrium and ventricle in our cases. We preferred the extra cardiac tunnel method using 16-20 size gore-tex tube grafts in all cases. The graft was anastomosed end-to-end to the right pulmonary artery, and to the vena cava inferior, 1 cm above the diaphragm. The graft anastomosis was performed end-to-end by transecting 1 cm above the diaphragm into the vena cava inferior. In the right pulmonary artery, it was performed medial to the anastomosis of the glans by the end-to-side method. In the right pulmonary artery, it was applied to the medial side by the end-side method. We routinely fenestrated a 5 mm diameter between the gore-tex graft and the right atrium. At the end of the operation, no graft deformation was observed and the oxygen saturation value increased over 95% (Figure 3). Our study was approved by the ethics committee decision dated 01.06.2021 and numbered 2021/11. The study was conducted in accordance with the Principles of the Declaration of Helsinki.

Results

Preoperative electrocardiogram (ECG) was sinus rhythm preoperatively in all patients. In echocardiographic findings, mean pulmonary artery pressures were 10(5-18) mmHg and mean pulmonary vascular resistance was 1.9 ± 0.7 Wood. The mean value of functional single ventricular ejection fraction before surgery was

59.4 ± 9.0%. Four patients had mild to moderate atrioventricular valve insufficiency, and three of them were in the under-4 age group. Other preoperative data of the patients are shown in table 2. Eleven patients were extubated on the first postoperative day and 6 patients were extubated within 2-3 days. All patients were given acetylsalicylic acid (100 mg/day) as antiaggregant prophylaxis after discharge. When we looked at the mortality rates of the cases, we lost 2 of 19 patients and the overall mortality rate was 10.5%. One of these cases was 38 months old and the other was 3 years old. We lost one of the patients as a result of acute heart failure that developed at the time of pump discharge, and the other due to low cardiac output during the intensive care period in the first 72 hours postoperatively. We used Extra Cardiac Membrane Oxygenation (ECMO) support in 3 patients. Supraventricular arrhythmia occurred in two patient in the early postoperative period. First of all, antiarrhythmic medical therapy was started. While one of these patients responded positively to medical treatment, the other could not be successful. Sinus rhythm was not achieved despite medical treatment. The cardioversion procedure that followed was successful, hemodynamically stable. However, we applied ICD implantation to this patient after 2 months due to intermittent extrasystolic beats. 7 patients were extubated within the first 24 hours and 8 patients within the first 48 hours. We encountered prolonged extubation in 2 patients, that is, in the first 3 days. While 13 patients could be discharged within 7-12 days, discharge in 4 patients varied between 15-35 days due to effusion due to prolonged chylothorax. There were no patients who needed hemodialysis with renal dysfunction. No mortality was observed in the near term after discharge. Outpatient follow-up of all patients was



Figure 1: Hypoplastic left pulmonary artery.




Figure 2: Stent placed in the right ventricular outflow tract and balloon dilatation of Glenn's anastomosis.




Figure 3: Our Fenestrated extracardiac type Fontan operation.

done at 3-month intervals. After discharge, the first echocardiography and other controls were performed 1 month later at 3-month intervals. The patients were followed for 1-3 years, with a mean of 17.2 ± 8.4 months. One year after the operation, signs of peripheral edema were observed in the areas around the eyes and in the lower extremities in 2 patients (Figure 4). Protein malnutrition was diagnosed in laboratory tests. In these patients, no significant gradients were detected in the anastomosis areas in the control angiography (Figure 5). One patient had partial graft thrombosis 3 months after discharge. The patient was hospitalized and a 1-week medical treatment was performed. The

Table 2: Preoperative baseline evaluation.

Ventricular morphology	
Dominant left ventricle	12
Dominant right ventricle	5
Biventricular	2
Preoperative EF	59.4 ± 9.0
Elevated creatinine	6
Dialysis	0
Pleural effusion	3
Systemic AVV regurgitation	4
Previous interventions (Branch PA/PDA stenting)	14
Cirrhosis	0
Prior stroke	0
Atrial arrhythmias	0
Permanent pacemaker	1

*EF: Ejection Fraction

*AVV: Atrioventricular Valve

*PA: Pulmonary Artery

*PDA: Patent Ductus Arteriosus



Figure 4: Edema findings around the eyes and in the body after 1 year in the patient who underwent Fontan surgery.

patient was discharged as the symptoms regressed. In addition, cardiac functions were seen well on echocardiography. In the postoperative 1-2 year echocardiographic follow-up of all patients, the functional single ventricle ejection fraction value was measured as $63.4 \pm 10\%$ ($p=0.085$). NYHA capacity of the patients was evaluated as 1 or 2. Postoperative data of the patients are shown in table 3.

Discussion

In 1971, the first fontan surgery was performed by Francis Fontan and Eugene Bauder. Although this surgery was originally described for



Figure 5: Control angiography 1 year after surgery, no stenosis in fontan anastomosis and fenestration spontaneously closed.

Table 3: Postoperative evaluation.

Discharge (In the 7-35 days)	In 7-35 days 17
Postoperative EF	% 63.4 ± 10 (p=0.085)
Cerebrovascular accident	0
Arrhythmias	2
Protein-losing enteropathy	2
Prolonged drainage (chylothorax)	4
Extubation (In the first 3 days)	17
Bleeding revision	1
Fontan graft thrombosis	1
Using ECMO	1 Successful; 1 Unsuccessful
ICD implantation	1
NHYA capacity	1-2

EF: Ejection Fraction

ECMO: Extra Cardiac Membrane Oxygenation

ICD: Intracardiac Defibrillator

NHYA: New York Heart Association

the treatment of tricuspid atresia, it was later chosen as the complementary stage of all univentricular pathologies [2]. In the 1970s, when it was first applied, mortality rates were between 20-35% [5]. In the 2000s, this rate decreased to around 7-10% with detailed research of some risk factors and various measurements. In the late 1970s, Choussat, et al. Published criteria that included the applicability of the Fontan procedure to which patients. After this period, the cases were evaluated according to the preoperative anatomical and physiological Fontan criteria. Decisions to wait or other palliative operations were made for cases that did not meet the criteria. One of the most important of these criteria is Pulmonary Vascular Resistance

(PVR). Normal PVR index should be <1.5 Wood unit. When the PVR index is >3 Wood units, the presence of pulmonary hypertension (PH) can be mentioned. In cases that are candidates for Fontan surgery, the desired limit is a maximum of 2-3 wood/unit [6]. Vasodilator application is sometimes recommended for patients with a basal PVR index between 4 and 6 Wood units. Some patients may have a 20% reduction in the PVR index with this. Another important factor is the mean Pulmonary Artery Pressure value (PAP). Normal value of PAP is 15-18 mmHg. In cases that are candidates for Fontan surgery, the average PAP should be <15 mmHg [7]. Normally developed right and left pulmonary arteries, functional single ventricle with good ejection fraction, absence of atrioventricular valve insufficiency, perfect venous return and normal sinus rhythm are other criteria for performing Fontan surgery [8]. The effect of age on mortality is also important. Likewise, both cases we lost were in the low age group. In the results of our clinic, the overall operative mortality rate can be seen as high as 10.5%. These two patients, especially under the age of 4, have contributed to this rate. On the other hand, there was no mortality in our patients over the age of 4 (17 patients). The mechanism by which the patient's age affects mortality is related to cardiovascular anatomy and pulmonary vascular structure reactivity that have not completed their development [9]. Among the causes of mortality in patients who have undergone Fontan surgery, arrhythmia, thromboembolism, cerebrovascular events and systemic venous return tract narrowing are also included [10]. As we know, there are 2 types of Fontan surgery as lateral tunnel and extra cardiac type. Both methods have different features. We prefer the extra cardiac type modification in all our cases. Considering that arrhythmia and narrowing of the venous return tract are more likely in the lateral type method, these risks are less in extra cardiac type fontan [11]. On the other hand, the extracardiac type is more advantageous in cases where the systemic venous connection is on the same side in the malposition of the apex of the heart. Using an excessively large size graft compared to the patient's IVC size may cause hemodynamic disturbance. We know that the IVC in an adult patient is approximately 20 mm. For children, this size is formed when they reach the age of about 15 kg and 3 years old. But sometimes, although body weight reaches the desired values, the development of cardiovascular structures is not the same in parallel. In this case, we think that it is correct to choose the conduit size based on the intraoperative evaluation of the actual IVC size. Our preference for the use of conduit was to apply one above the IVC size that we measured intra-operatively, especially in 3-year-old cases. Generally, there is an opinion that 16-18 mm conduits are ideal for 3-3,5 year old children [12]. Lardo AC, et al. Alexi-Meskishvili VA, et al. have important ideas in their conduit-IVC size selection. In cases where 20 mm and larger conduits were used at this age, stagnation in excess fluid volume in the lateral graft and backflow in the lateral part during expiration was shown. There are still various controversial opinions regarding the opposition or shifting of the superior vena cava with conduit anastomosis in cases with Glenn shunt. Our preference is to place the conduit anastomosis in a more medial area. With this method, we consider a more even and better distribution of blood to the right and left lungs [13,14]. Some patients may have difficulties in keeping up with the new physiology after the fontan operation. The most important of these is prolonged pleural effusions. Especially chylothorax, protein-losing enteropathy and some lung problems. We discharged a total of 4 patients late due to prolonged chylothorax consistency drainage. Appropriate diet was used in the treatment of chylothorax. We supported patients' intake of adequate calories and protein while restricting their consumption of fat containing medium-chain triglycerides [15]. Chronic passive stasis may occur in the venous system and liver after Fontan surgery. Secondary inflammation

and increased platelet activity and thromboembolic complications may occur. This can be considered an unwanted disadvantage [16]. Stroke is one of the most important complications in patients who have undergone Fontan surgery [17]. We have not encountered such events in our series. Anticoagulation and antiaggregation use in different combinations is required against thromboembolism. In all patients, we routinely preferred low molecular weight heparin for the first 48 hours postoperatively and then aspirin tablet 100 mg/day for 6 months. We did not encounter any thrombosis event in the short and medium term impression. In this regard, the choice of treatment protocol is subjectively different in separate centers. After thromboembolic events develop, the mortality rate can be 38% despite aggressive treatment [18]. When we compared the functional single ventricle load before and after surgery, it was more spherical, the muscle mass was partially hypertrophic and the volume was higher before the operation. Positive changes were reflected in these parameters after the operation. The diastolic functions of the ventricle are important, especially in providing pulmonary circulation. Sometimes, after the fontan procedure, ventricular diastolic function deteriorates secondary to relaxation disorder, and this slows down the systemic venous circulation [19]. Throughout our study, we did not follow any patient with ventricular systolic and diastolic dysfunction, except for 2 patients we lost. Previously known myocardial dysfunction or atrioventricular valve insufficiency was not observed in the patients who were lost. In a 10-year study by Mira Fauziah et al. [20] systemic ventricular morphology was not associated with the 10-year survival rate after Fontan surgery. The probable cause was the problem of meeting the systemic venous return by the pulmonary system, diameter mismatch or increased pulmonary vascular resistance. Today, Fontan surgery is usually performed by applying fenestration [21]. In our case series, we applied 3-5 mm of routine fenestration in all cases. Fenestration slightly increases cardiopulmonary bypass time, saturation value may be slightly lower in the early period, but decreases the incidence of arrhythmia [22]. We believe that fenestration reduces the incidence of postoperative effusion in the short term and shortens the length of hospital stay. Fenestration helps reduce the pressure of the inferior vena cava. It usually closes on its own in the long term. In the angiography we performed for 1 year sonar control, we saw that the fenestration was closed (Figure 5). Some centers also perform Fontan surgery without the use of fenestration. In most studies, approximately 70% of these fenestrations closed spontaneously. Fontan surgery can be performed easily as long as it meets all the surgical indication conditions mentioned above. Here, along with other indications, the main goal is the systemic venous circulation to be included in the pulmonary system without interfering with the heart and to ensure circulation [23-26]. This procedure has the hemodynamic advantages mentioned above and is still a good palliative operation for single ventricle cases.

Conclusion

In order to reduce mortality and morbidity, the result we have inferred from our case series is to move the age factor to more advanced ages as possible. We have the opportunity to say this, especially since the patients we lost are under the age of 4. If the preoperative oxygen saturation value of the patients is not critical, if the functional single ventricle function is normal, if there is no significant leakage in the atrioventricular valve, if the physical exertion capacity and hemodynamic circulation findings are close to normal, it is useful to wait as long as possible between the Glenn shunt and Fontan surgery. In conclusion, we think that the morbidity and mortality risk of surgery performed at an older age is low.

Conflict of Interests

There is no conflict of interest in the preparation and publication of this article.

Funding

There is no financial support for the research and authorship process of this article.

Author Contributions

All authors read and approved the final version of the manuscript.

References

1. Wortmann LG, Dragulescu A, Drolet C, Chaturvedi R, Kotani Y, et al. (2013) Determinants and clinical significance of flow via the fenestration in the Fontan pathway: a multimodality study. *Int J Cardiol* 168: 811-817.
2. Aboul Hosn JA, Shavelle DM, Castellon Y, Criley JM, Plunkett M, et al. (2007) Fontan operation and the single ventricle. *Congenit Heart Dis* 2: 2-11.
3. Lee C, Lee CH, Hwang SW, Lim HG, Kim SJ, et al. (2007) Midterm follow-up of the status of Gore-Tex graft after extracardiac conduit Fontan procedure. *Eur J Cardiothorac Surg* 31: 1008-1012.
4. Bouhout I, Ali WB, Khalaf D, Raboisson MJ, Poirier N (2020) Effect of Fenestration on Fontan Procedure Outcomes: A Meta-Analysis and Review. *Ann Thorac Surg* 109: 1467-1474.
5. Corno AF, Owen M, Cangiani A, Hall EJC, Rona A (2019) Physiological Fontan Procedure. *Front Pediatr* 7: 196.
6. Egbe AC, Connolly HM, Miranda WR, Ammash NM, Hagler DJ, et al. (2017) Hemodynamics of Fontan Failure: The Role of Pulmonary Vascular Disease. *Circ Heart Fail* 10: e004515.
7. Vasdev S, Chauhan S, Talwar S, Pandey A, Kiran U, et al. (2013) Monitoring pulmonary artery pressure in children undergoing the Fontan procedure. *Asian Cardiovasc Thorac Ann* 21: 303-305.
8. Ohuchi H (2017) Where Is the "Optimal" Fontan Hemodynamics? *Korean Circ J* 47: 842-857.
9. Akintoye E, Veldtman GR, Miranda WR, Connolly HM, Egbe AC (2019) Optimum age for performing Fontan operation in patients with univentricular heart. *Congenit Heart Dis* 14: 138-139.
10. VanHise K, Li L, Smith E, Condon N (2018) Embolic stroke in a postpartum patient with Fontan physiology. *BMJ Case Rep* 2018: bcr2017222684.
11. Kwak JG, Lee JR, Jang WS, Bae EJ (2008) Extracardiac Fontan with T-shape conduit in non-confluent pulmonary arteries. *J Cardiothorac Surg* 3: 7.
12. Quinones JA, Deleon SY, Bell TJ, Cetta F, Moffa SM, et al. (1997) Fenestrated fontan procedure: evolution of technique and occurrence of paradoxical embolism. *Pediatr Cardiol* 18: 218-221.
13. Lardo AC, Webber SA, Friebs I, Nido PJD, Cape EG (1999) Fluid dynamic comparison of intra-atrial and extracardiac total cavopulmonary connections. *J Thorac Cardiovasc Surg* 117: 697-704.
14. Meskishvili VA, Ovroutski S, Ewert P, Dähnert I, Berger F, et al. (2000) Optimal conduit size for extracardiac Fontan operation. *Eur J Cardiothorac Surg* 18: 690-695.
15. Johnson JN, Driscoll DJ, O'Leary PW (2012) Protein-losing enteropathy and the Fontan operation. *Nutr Clin Pract* 27: 375-384.

16. Keung CY, Zentner D, Gibson RN, Phan DKH, Grigg LE, et al. (2020) Fontan-associated liver disease: pathophysiology, investigations, predictors of severity and management. *Eur J Gastroenterol Hepatol* 32: 907-915.
17. Firdouse M, Agarwal A, Chan AK, Mondal T (2014) Thrombosis and thromboembolic complications in fontan patients: a literature review. *Clin Appl Thromb Hemost* 20: 484-492.
18. John AS (2015) Fontan Repair of Single Ventricle Physiology: Consequences of a Unique Physiology and Possible Treatment Options. *Cardiol Clin* 33: 559-569.
19. Goldstein BH, Connor CE, Gooding L, Rocchini AP (2010) Relation of systemic venous return, pulmonary vascular resistance, and diastolic dysfunction to exercise capacity in patients with single ventricle receiving fontan palliation. *Am J Cardiol* 105: 1169-1175.
20. Fauziah M, Lilyasari O, Liastuti LD, Rahmat B (2018) Systemic ventricle morphology impact on ten-year survival after Fontan surgery. *Asian Cardiovasc Thorac Ann* 26: 677-684.
21. Li D, Li M, Zhou X, An Q (2019) Comparison of the fenestrated and non-fenestrated Fontan procedures: A meta-analysis. *Medicine (Baltimore)* 98: e16554.
22. Mery CM, León LED, Diaz DT, Ocampo EC, Dickerson HA, et al. (2019) Contemporary Outcomes of the Fontan Operation: A Large Single-Institution Cohort. *Ann Thorac Surg* 108: 1439-1446.
23. Lemler MS, Scott WA, Leonard SR, Stromberg D, Ramaciotti C (2002) Fenestration improves clinical outcome of the fontan procedure: a prospective, randomized study. *Circulation* 105: 207-212.
24. Patel ND, Friedman C, Herrington C, Wood JC, Cheng AL (2021) Progression in Fontan conduit stenosis and hemodynamic impact during childhood and adolescence. *J Thorac Cardiovasc Surg* 162: 372-380.
25. Wan D, Grewal J, Barlow A, Kiess M, Human D, et al. (2020) Atrial arrhythmias and thromboembolic complications in adults post Fontan surgery. *Open Heart* 7: e001224.
26. AlZahrani A, Rathod R, Krimly A, Salam Y, AlMarzoog AT, et al. (2020) The Adult Patient with a Fontan. *Cardiol Clin* 38: 379-401.